

540,196

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 July 2004 (08.07.2004)

PCT

(10) International Publication Number
WO 2004/056202 A1

(51) International Patent Classification⁷: **A23L 1/22**

(21) International Application Number:
PCT/IB2002/005529

(22) International Filing Date:
20 December 2002 (20.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): **COUNCIL OF SCIENTIFIC AND INDUSTRIAL RESEARCH [IN/IN];** Rafi Marg, 110 001 New Delhi (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SRINIVAS, Pulabhatla [IN/IN];** Central Food and Technological Research Institute, Mysore, 570 013 Karnataka (IN). **SULOCHANAMMA, Guruguntla [IN/IN];** Central Food and Technological Research Institute, Mysore, 570 013 Karnataka (IN). **RAGHAVAN, Bashyam [IN/IN];** Central Food and Technological Research Institute, Mysore, 570 013 Karnataka (IN). **GURUDUTT, Kambadoor, Nagarajarao [IN/IN];** Central Food and Technological Research Institute, Mysore, 570 013 Karnataka (IN).

(74) Agent: **GABRIEL, Devados, Calab; K & S Partner, 84-C, C6 Lane, Off Central Avenue, Sainik Farms, 110 062 New Delhi (IN).**

(81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,**

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)**

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **AN IMPROVED PROCESS FOR THE STABILIZATION OF 2-ACETYL-1-PYRROLINE, THE BASMATI RICE FLAVOURANT**

(57) Abstract: The present invention provides an improved process for the stabilization of 2-acetyl-1-pyrroline, a basmati aroma producing principle, the said process comprising steps of: dissolving a binder in water containing few drops of an emulsifier, adding an ethanol solution of 2-acetyl-1-pyrroline, homogenizing for a time period of 3 to 5 minutes, and drying the solution to obtain the stabilized flavor 2-acetyl-1-pyrroline in a dispersible dry powder form.

WO 2004/056202 A1

AN IMPROVED PROCESS FOR THE STABILIZATION OF 2-ACETYL-1-PYRROLINE, THE BASMATI RICE FLAVOURANT

Technical field

The present invention relates to 'An improved process for stabilization of 2-acetyl-1-pyrroline, the principle aroma component of basmati and other varieties of scented rice and also of processed cereal and grain products'.

The main uses of the present invention are (i) making the basmati rice flavouring in easily dispersible powder form and (ii) imparting stability to an otherwise labile aroma chemical. These products are of value in the art of application of flavor to foodstuffs like rice and bakery products and related preparations.

Background and prior art references

Scented rice is one of the varieties of rice¹⁻⁶ having a characteristic, strong aroma when cooked. Varieties of Basmati in Southeastern Asia, Della in America, Milagrosa in Philippines, 'Khao Dawk Mali 105' in Thailand, 'Seratus Malam' in Indonesia and Heiri in Japan are the major scented rice varieties. In certain parts of the world where these varieties are cultivated, scented rice is highly desired and sometimes used for special purposes.

2-acetyl - 1-pyrroline was first identified in rice by Buttery et al⁷⁻⁸ in 1982. Its odor is described as that of cooked rice and popcorn-like. A method of its synthesis is described starting from 2-acetylpyrrole⁹. A method for its quantitative analysis has been developed by Buttery *et al*¹⁰ and later refined by Tanchotikul et al¹¹. This compound is found in all scented rice varieties in various quantities, which is more than that found in non-scented varieties of rice (table 1).

Mol. formula: C₆H₉ON;
Mol. Wt. 111;
CAS Registry No: 99583-29-6;
Structural Formula




Table 1: LEVEL OF 2-ACETYL-1-PYRROLINE IN SCENTED RICE VARIETIES⁶

Rice sample	ppb
Malagkit Sungsong (brown)	760
Basmati 370 (brown)	610
IR 841-76-1 (brown)	560
Texas long grain (Bluebell var; polished)	60
Khao Dawk Mali 105 (brown)	710
Azucena (brown)	570
Hieri (brown)	360

5 2-acetyl - 1-pyrroline can be characterized by a comparison of its retention time and MS data with those of a reference standard or reported in literature. The mass spectrum of 2-acetyl - 1-pyrroline has major peaks at m/z 111 (M^+ , 5 % abundance, 83 (11), 69 (11), 68 (8), 67 (0.2), 55(2), 52 (0.9), 54 (0.2) and 43 (100), 42 (24) and 41 (50). It's Kovats index is 1320 on a Pyrex glass capillary column coated with Carbowax 20M. It's
 10 IR spectrum in CCl_4 displays absorption maxima at 1695, 1620, 1435, 1370, 1340, 1250, 1080, 1000, 975 and 940 cm^{-1} .

The compound is a colorless liquid when freshly prepared and purified. It needs to be immediately protected from light and air and preserved in sealed vials under vacuum at temperatures below -20°C . Even at under these conditions it is reported to turn to red on
 15 storage and eventually become darker on longer storage. A conjugated polymer, resulting from conjugation of the carbonyl groups with the 5-position of other molecules resulting in a polymeric product, is believed to be the cause of its instability. For this reason the compound is more stable and better preserved in dilute, especially aqueous, solutions wherein it is stable for several months at low temperatures ($<-20^\circ\text{C}$).

20 It seems reasonable that 2-acetyl-1-pyrroline has a similar origin to the bread aroma compound 2-acetyl-1,4,5,6-tetrahydropyridine¹². It is assumed that the intermediate formed after decarboxylation of proline residue, is hydrolyzed with the formation of 1-pyrroline. 2-acetyl-1-pyrroline results from the acetylation 1-pyrroline by 2-oxopropanal. Many syntheses of 2-acetyl-1-pyrroline are reported in recent literature¹³⁻¹⁷.

25 Reference may be made to the patent United States Patent (Richard; Travis, 6,274,183, 2001), wherein a food coating composition is manufactured from fragrant or

scented rices (aromatic rices) by grinding the fragrant rice having a 2-acetyl-1-pyrroline concentration of at least 40 ppb by weight to a predetermined particle size at a predetermined temperature to obtain a food coating composition with uniform adherence to the food product, as well as improved crispness. The resulting food coating composition is claimed to provide a unique and distinct aroma and taste, which is derived from 2-acetyl-1-pyrroline. The main drawback of this is the low levels of the flavor that would be available on dilution in the final product.

Reference may be made to the United States Patents (Duby; Philippe (Prilly, CH); Huynh-Ba; Tuong (Pully, CH) 5,280,127, 1994; Duby; Philippe ; Huynh-Ba; Tuong , 5,401,521 & 5,446,171, 1995 and Duby; Philippe ; Huynh-Ba; Tuong , 5,512,290, 1996), wherein a powder-form composition of 2-acetyl-1-pyrroline incorporated with a support of maltodextrin and/or cyclodextrin. The composition is prepared by hydrolyzing a 2-(1-alkoxyethenyl)-1-pyrroline with an acid to obtain a reaction medium, adding an equi-molar amount of a base to the reaction medium to obtain a neutralized reaction medium containing 2-acetyl-1-pyrroline, combining maltodextrin and/or cyclodextrin with the neutral reaction medium to obtain a support solution and freeze-drying the support solution to obtain the composition. The main drawback of these methods is that the flavor needs to be released from the precursor by the neutralization with a base and the resultant salt would remain in the product.

Reference may be made to another United States Patent (Buttery; Ronald G.; Ling; Louisa C.; Juliano; Bienvenido O., 4,522,838, 1985), wherein the compound, 2-acetyl-1-pyrroline and its use in flavoring foods, particularly in imparting a scented rice flavor to foods is disclosed. The drawback of this invention is that the flavorant is used in its salt form and therefore needs to be released prior to use.

Objective of the invention

The main objective of the present invention is to provide an improved process for stabilization of 2-acetyl-1-pyrroline, a principle basmati aroma component', which obviates the aforesaid drawbacks of the processes described in the prior art.

Summary of the Invention

Accordingly, the present invention provides, 'an improved process for stabilization of 2-acetyl-1-pyrroline, the basmati rice flavourant', which comprises synthesis of 2-

acetyl-1-pyrroline by a known method, incorporation of the flavor in a binding material like a gum or starch from a vegetable source in water emulsion using an emulsifier followed by drying the mixture by either vacuum shelf drying at temperatures of 30-60°C and reduced pressure of 24'' or spray drying at inlet air temp. 140° C, outlet temp. 80° C and a feed rate of 80 ml / min and obtaining the flavor in an easily dispersible dry powder form for flavoring rice and related products.

Detailed Description of the Invention

Accordingly, the present invention provides an improved process for the stabilization of 2-acetyl-1-pyrroline, a basmati aroma producing principle, the said process comprising steps of:

- a) dissolving a binder in water containing few drops of an emulsifier,
- b) adding an ethanol solution of 2-acetyl-1-pyrroline to step (a) solution,
- c) homogenizing step (b) solution for a time period of 3 to 5 minutes, and
- d) drying the homogenized solution of step (c) to obtain the stabilized flavor 2-acetyl-1-pyrroline in a dispersible dry powder form.

In an embodiment of the invention, the binder used in step (a) is from a vegetable source and which is selected from a group consisting of gum acacia, starch or mixtures thereof.

Still another embodiment of the invention, the ratio of 2-acetyl-1-pyrroline and the binder used is in the ratio of 0.1:2000 to 1.0: 2000.

Still The process of claim 1, wherein in step (a), the emulsifier used is selected from a group consisting of Tween 80, Tween 60 and more preferably Tween 60.

Yet another embodiment, 2- acetyl-1-pyrroline used is prepared by adopting known methods.

In yet another embodiment drying in step (d) is carried out by vacuum shelf drying or spray drying.

Yet another embodiment, the vacuum shelf drying is performed at reduced pressure of 24'' and at a temperature in the range of 30° - 60° C, and spray drying is carried by using a feed rate of 80 ml/min. with an inlet air temperature of 140° C and outlet temperature of 80° C.

One more embodiment of the invention provides a process, wherein the stabilized flavor obtained is used for flavoring rice and related products.

The following examples are given by way of illustration of the present invention and therefore should not be construed to limit the scope of the present invention.

EXAMPLE 1:

In a typical experiment gum acacia (200 g) was dissolved in 2 l water, a solution of 100 mg of 2-acetyl-1-pyrroline (> 95 % purity, synthetic sample) in ethanol (3 ml) added, the mixture homogenized for 3 min. and spray-dried under the following conditions: Inlet air temp: 140°C; outlet temp: 80°C; feed rate: 80 ml / min. The yield was 90 %.

Preparation of Sweetened basmati rice**A) Rice (added basmati flavor)**

Rice (100 g) was taken, added water (1:3) and cooked for 15 mins. Cooked rice was macerated in a mixer for 1 min. Condensed milk was diluted with water (1:1), and the diluted milk (150 ml) and encapsulated basmati flavor (1% w /w basis on raw rice) were added into the cooked rice and chilled.

B) Natural basmati rice

Non-scented rice (100 g) was taken, added water (1:3) and cooked for 15 mins. Cooked rice was macerated in a mixer for 1 min. The condensed milk diluted 1:1 (150 ml) was added to the cooked rice and chilled.

Evaluation of sweetened Basmati rice

Rice with added basmati flavor and natural basmati rice were evaluated by sensory analysis on a 10-point scale for quality attributes such as aroma, texture and overall acceptability by a panel of 10 judges. The results reveal that the rice with added flavor is preferred and liked more in comparison to the natural basmati rice with respect to the aroma and overall quality.

EXAMPLE 2

In a typical experiment binding material (see table below) was dissolved in 2 l water, Tween – 60 (4 drops) and a solution of 2-acetyl-1-pyrroline (> 95 % purity, synthetic sample) in ethanol (3 ml) added, homogenized for 5 min. and vacuum shelf-dried under the following conditions. 30-60°C / 24''. The results are tabulated below.

Binding material	Quantity (g)	2-AP (mg)	Yield of Encapsulated Product (g)
Gum acacia	200	100	143
Starch (HICAP)	200	45	89

Gum acacia + Starch (HICAP)	100 + 100	45	98
--------------------------------	-----------	----	----

EXAMPLE 3

Also in another experiment starch (200 g) was dissolved in 2 l water, Tween (4 drops) and a solution of 110 mg of 2-acetyl-1-pyrroline (> 95 % purity, synthetic sample) in ethanol (3 ml) added and the mixture homogenized for 5 min. and spray-dried under the following conditions. Inlet-air temp. 140⁰ C, outlet temp. 80⁰ C and a feed rate of 80 ml / min. The yield was 162.2 g.

10 Incorporation of synthetic basmati flavor in to rice and its comparison with Natural basmati rice

Preparation of Sweetened basmati rice

A) Rice (added basmati flavor)

Rice (100 g) was taken, added water (1:3) and cooked for 15 mins. Cooked rice was macerated in a mixer for 1 min. Condensed milk diluted with water (1:1, 150 ml) and encapsulated basmati flavor (1% w /w basis on raw rice) were added into the cooked rice and chilled.

B) Natural basmati rice

Rice (100 g) was taken, added water (1:3) and cooked for 15 mins. Cooked rice was macerated in a mixer for 1 min. The diluted condensed milk (150 ml) was added to the cooked rice and chilled.

Evaluation of sweetened basmati rice

Rice with added basmati flavor and natural basmati rice were evaluated by sensory analysis on a 10-point scale for quality attributes such as aroma, texture and overall acceptability by a panel of 10 judges. The results revealed that the rice with added flavor was preferred and liked more in comparison to the natural basmati rice with respect to the aroma and overall quality. However the texture was described as "pasty" in both the samples.

30 REFERENCES

1. *Volatile Compounds in Foods and beverages*, Maarse H, Marcel Dekker, New York, 1991, pp79-89.
2. Buttery R G, Turnbaugh J G and Ling L C, Contribution of volatiles to rice aroma, *J Agric Food Chem.*, 1988, **36(5)**, 1006-1009
3. Lin C F, Hsieh, T C.-Y and Hoff B J, Identification and quantification of the "pop-

4. Seitz L M, Wright R L, Waniska R D and Rooney L W, Contribution of 2-acetyl pyrroline to odors from wetted ground pearl millet, *J Agric Food Chem*, 1993, 41(6), 955-958.
5. Widjaja R, Craske J D and Wootton M, Comparative studies on volatile components of non-fragrant and fragrant rices, *J Sci Food Agric*, 1996, 70, 151-161.
6. Paule C M and Powers J J, Sensory and chemical examination of aromatic and non-aromatic rices, *J Food Sci.*, 1989, 54(2), 343-346.
7. Buttery R G, Ling L C and Juliano B O, 2-acetyl-1-pyrroline : an important aroma component of cooked rice, *Chem and Ind*, 1982, 958-959
- 10 8. Buttery R G, Juliano B O and Ling L C, Identification of rice aroma compound 2-acetyl-1-pyrroline in pandana leaves, *Chem and Ind*, 1983, 478
9. Buttery R G, Ling L C, Juliano B G and Turnbaugh J G, Cooked rice aroma and 2-acetyl pyrroline, *J Agric Food Chem*, 1983, 31(4), 823-826.
- 15 10. Buttery R G, Ling L C and Mon T R, Quantitative analysis of 2-acetyl pyrroline, *J Agric Food Chem*, 1986, 34(1), 112-114
11. Tanchotikul U and Hsieh T C.-Y, An improved method for quantification of 2-acetyl-1-pyrroline, a "popcorn"-like aroma, in aromatic rice by high-resolution gas chromatography / mass spectrometry / selected ion monitoring, *J Agric Food Chem*, 1991, 39 (5), 944-947.
- 20 12. Schieberle P and Grosch W, Identification of the volatile flavour compounds of wheat bread crust - comparison with rye bread crust, *Zeitschrift fur Lebensmittel untersuchung and Forchung*, 1985, 180(6), 474-478.
13. De Kimpe N, Stevens C V and Keppens M, Synthesis of 2-acetyl pyrroline, the principle rice flavour component, *J Agric Food Chem*, 1993, 41(9), 1458-1461.
- 25 14. De Kimpe N and Keppens M, Novel syntheses of the major flavour components of bread and cooked rice, *J Agric Food Chem*, 1996, 44(6), 1515-1519
15. Hofmann T and Schieberle P, New and convergent synthesis of the important roasty, pop-corn like smelling food aroma compounds, 2-acetyl-1-pyrroline and 2-acetyltetrahydropyridine from their corresponding cyclic α -amino acids, *J Agric Food*
- 30 *Chem*, 1998, 46(6), 2270-2277
16. Hofmann T and Schieberle P, Penicillin acylase-mediated synthesis of 2-acetyl-1-pyrroline and 2-propionyl-1-pyrroline, key roast smelling odourant in Food. Inclusion complexes with β -cyclodextrin and their NMR & MS characterisation, *J Agric Food Chem.*, 1998, 46(6), 616-619.
- 35 17. Favino T F, Fronza G, Fuganti C, Fuganti D, Grasselli and Mele A, 2-oxopropanol, hydroxy-2-propanone and 1-pyrroline - important intermediates in the generation of the roast-smelling food flavour compounds, 2-acetyl-1-pyrroline and 2

Claims

1. An improved process for the stabilization of 2-acetyl-1-pyrroline, a basmati aroma producing principle, the said process comprising steps of:
 - a) dissolving a binder in water containing few drops of an emulsifier,
 - 5 b) adding an ethanol solution of 2-acetyl-1-pyrroline to step (a) solution,
 - c) homogenizing step (b) solution for a time period of 3 to 5 minutes, and
 - d) drying the homogenized solution of step (c) to obtain the stabilized flavor 2-acetyl-1-pyrroline in a dispersible dry powder form.
2. The process of claim 1, wherein in step (a), the binder used is from a vegetable source.
- 10 3. The process of claim 2, wherein the binder used is selected from a group consisting of gum acacia, starch or mixtures thereof.
4. The process of claim 1, wherein the ratio of 2-acetyl-1-pyrroline and the binder used is in the ratio of 0.1 to 1.0: 2000.
5. The process of claim 1, wherein in step (a), the emulsifier used is selected from a group
15 consisting of Tween 80, Tween 60 and more preferably Tween 60.
6. The process of claim 1, wherein 2- acetyl-1-pyrroline used is prepared by adopting known methods.
7. A process of claim 1, wherein in step (d) the drying is performed by vacuum shelf drying or spray drying.
- 20 8. A process of claim 7, wherein vacuum shelf drying is performed at reduced pressure of 24'' and at a temperature in the range of 30° - 60° C.
9. A process of claim 7, wherein spray drying is carried by using a feed rate of 80 ml/min. with an inlet air temperature of 140° C and outlet temperature of 80° C.
10. A process of claim 1, wherein the stabilized flavor obtained is used for flavouring rice
25 and related products.

INTERNATIONAL SEARCH REPORT

In **onal Application No**
PCT/IB 02/05529

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A23L1/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, FSTA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 512 290 A (DUBY PHILIPPE ET AL) 30 April 1996 (1996-04-30) cited in the application column 1, line 40 - line 55 column 6; example 6	1-10
A	US 6 274 183 B1 (RICHARD TRAVIS) 14 August 2001 (2001-08-14) cited in the application the whole document	1-10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

7 August 2003

Date of mailing of the international search report

22/08/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Inceisa, L

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 02/05529

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5512290	A	30-04-1996	CH 683995 A5	30-06-1994
			US 5401521 A	28-03-1995
			AT 190050 T	15-03-2000
			AU 659888 B2	01-06-1995
			AU 2851592 A	03-06-1993
			CA 2084256 A1	03-06-1993
			DE 69230717 D1	06-04-2000
			DE 69230717 T2	17-08-2000
			EP 0545085 A1	09-06-1993
			JP 2562267 B2	11-12-1996
			JP 5262725 A	12-10-1993
			NO 924483 A ,B,	03-06-1993
			NO 954994 A ,B,	08-12-1995
			NZ 245259 A	28-08-1995
			NZ 270635 A	28-08-1995
			US 5446171 A	29-08-1995
			US 5280127 A	18-01-1994
			ZA 9208956 A	21-02-1994
US 6274183	B1	14-08-2001	NONE	